



20

(Knowledge for Development)

KIBABII UNIVERSITY
UNIVERSITY EXAMINATIONS
2016/2017 ACADEMIC YEAR
SECOND YEAR FIRST SEMESTER
SPECIAL/ SUPPLEMENTARY EXAMINATION
FOR THE DEGREE OF BACHELOR OF SCIENCE (IT)

COURSE CODE: MAT 211

COURSE TITLE: LINEAR ALGEBRA

DATE: 15/09/17

TIME: 11.30 AM -1.30 PM

INSTRUCTIONS TO CANDIDATES

Answer Question One and Any other TWO Questions

TIME: 2 Hours

This Paper Consists of 4 Printed Pages. Please Turn Over.

QUESTION ONE (30 MKS)

(a) (i) Define a vector space over \mathbb{R} . (3mks)

(ii) Show that an n –dimensional Euclidean space is a vector space (4mks)

(b) Show that the function $h(x) = 4x^2 + 3x - 7$ lies in the space $span\{f, g\}$ generated by $f(x) = 2x^2 - 5$ and $g(x) = x + 1$. (3mks)

(c) (i) Distinguish between linear dependence and independence. (1mks)

(ii) Determine whether the set $\{(1,2,0), (0,1, -1), (1,1,2)\}$

is linearly dependent in \mathbb{R}^3 . (2mks)

(d) Prove that $\{(1,3, -1), (2,1,0), (4,2,1)\}$ is a basis for \mathbb{R}^3 . (3mks)

(e) (i) Show that the row space and the column space of a matrix A have the same dimension. (3mks)

(ii) Find the basis for the row space of the following matrix $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 5 & 4 \\ 1 & 1 & 5 \end{pmatrix}$ and determine its rank. (3mks)

(f) (i) Define a linear transformation (1mks)

ii) Prove that the following transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(x, y) = (2x, x + y)$ is linear. (2mks)

(g) Define

i) Kernel

ii) Range (2mks)

(h) Find the kernel and the Range of the linear operator: $T(x, y, z) = (x, y, 0)$. (3mks)

QUESTION TWO (20MKS)

(a) Determine the kernel and the Range of the transformation defined by the

matrix = $\begin{pmatrix} 1 & 2 & 3 \\ 0 & -1 & 1 \\ 1 & 1 & 4 \end{pmatrix}$. (6mks)

(b) Given $T: U \rightarrow V$ is a linear transformation, show that

$$\dim \text{Ker}(T) + \dim \text{range}(T) = \dim \text{domain } T. \quad (7\text{mks})$$

(c) Let T be a linear transformation of $\mathbb{R}^n \rightarrow \mathbb{R}^n$. Show that T is invertible if and only if it is singular, its inverse is unique and it is linear. (7mks)

QUESTION THREE (20MKS)

(a) Define an invertible linear transformation on \mathbb{R}^n . (2mks)

(b) Show that the linear transformation T is one-to-one if and only if the kernel is a zero vector. (4mks)

(c) Consider the linear transformation: $T(x, y, z) = (3x + 4y, 5x + 7y)$ of $\mathbb{R}^2 \rightarrow \mathbb{R}^2$.

(i) Prove that T is invertible and find the inverse of T . (4mks)

(ii) Determine the pre-image of the vector $(1, 2)$. (2mks)

(d) i) Define a vector subspace (2mk)

ii) Show that the set U of a 2×2 diagonal matrices is a subspace of the vector space M_{22} of 2×2 matrices. (3mks)

iii) Let W be a set of vectors of the form $(a, a, a + 2)$.

Show that W is not a subspace of \mathbb{R}^3 . (3mks)

QUESTION FOUR (20 MKS)

(a) i) Define a vector basis. (2mks)

ii) Let $\{v_1, \dots, v_n\}$ be a basis for a vector space V . If $\{w_1, \dots, w_n\}$ is a set of more than n vectors in V . Show that this set is linearly dependent. (5mks)

(b) Prove the following

i) the origin is a subspace of \mathbb{R}^3 .

ii) the one dimensional subspaces of \mathbb{R}^3 are lines through the origin.

iii) the two dimensional subspaces of \mathbb{R}^3 are planes through the origin. (9mks)

c) Let $\mathbb{R}^n \rightarrow \mathbb{R}^n$ be defined by $T(u) = A(u)$. Show that the range of T is spanned by the column vector of A . (4mks)

QUESTION FIVE (20MKS)

(a) Solve the following homogeneous system of linear equations. Interpret the set of solutions as a subspace and sketch the subspace of solutions.

$$x_1 + 2x_2 + 3x_3 = 0$$

$$-x_2 + x_3 = 0$$

$$x_1 + x_2 + 4x_3 = 0$$

(10mks)

(b) Analyse the solutions to the following system of equations

$$x_1 - 2x_2 + 3x_3 + x_4 = 1 \quad x_2 + 2x_3 - x_4 = 4$$

$$3x_1 - 5x_2 + 5x_3 = 5$$

$$x_1 - x_2 - 2x_4 = 3$$

(10mks)