

Handwritten signature or mark in red ink.



# KIBABII UNIVERSITY

**UNIVERSITY EXAMINATIONS  
2020/2021 ACADEMIC YEAR**

**FOURTH YEAR SECOND SEMESTER  
MAIN EXAMINATIONS**

**FOR THE DEGREE OF B.ED (SCIENCE)**

**COURSE CODE: SCH 440:**

**COURSE TITLE: ELECTROCHEMISTRY**

**DURATION: 2 HOURS**

**DATE: 15/07/2021**

**TIME: 2:00-4:00PM**

---

## **INSTRUCTIONS TO CANDIDATES**

- Answer **QUESTION ONE** (Compulsory) and any other two (2) Questions.
- Indicate **answered questions** on the front cover.
- Start every question on a new page and make sure question's number is written on each page.

This paper consists of 6 printed pages. Please Turn Over



**KIBU observes ZERO tolerance to examination cheating**


**Question One (30 Marks)**

- a)
- i. The Lead-acid cell, also called an acid accumulator has the overall spontaneous reaction below.  


$$\text{Pb}_{(s)} + \text{PbO}_{2(s)} \rightarrow \text{PbSO}_{4(s)}$$
    - i. Write the equation of the reaction at the **anode**. [02]
    - ii. Write the equation of the reaction at the **Cathode**. [02]
    - iii. Differentiate between a galvanic cell and an electrolytic cell. [02]
- b) An element X has r.a.m of 88. When a current of 0.5A was passed through the fused chloride of X for 32minutes and 10 seconds, 0.44g of X was deposited at the cathode. (Use 1 Farad = 96,500 coulombs)
- i. Calculate the quantity of electricity needed to liberate one mole of X. [04]
  - ii. Write the formular for the cation of X. [01]
  - iii. Write formular for the chloride of X. [01]

- c)
- i) For a galvanic cell combining Zn and Cu, calculate the standard cell potential  $E^\circ$  ( given standard reduction potential for  $\text{Zn}^{2+}$  is -0.76V and that for  $\text{Cu}^{2+}$  is +0.34V) [03]
  - ii) Calculate the cell potential for the Zn//Cu cell at  $[\text{Zn}^{2+}_{(aq)}] = 10\text{M}$  and  $[\text{Cu}^{2+}_{(aq)}] = 0.1\text{M}$  [03]

- d) The oxidation and reduction half cell reactions of the following overall process exist in separate half cells.



Given the stoichiometric equation is,



[05]

Species	Concentration
$\text{Cr}^{3+}_{(aq)}$	$2.0 \times 10^{-3}$
$\text{Cr}_2\text{O}_7^{2-}_{(aq)}$	2.0
$\text{H}^+_{(aq)}$	1.0
$\text{I}_{(aq)}$	1.0

- e) Dichromate (VI) ions are powerful oxidising agents and are reduced to chromium III ions. This colour change was once used in 'breath test' apparatus to determine if a driver had consumed excessive alcohol. Explain the term 'redox reaction' [01]

f) Aluminium metal is extracted from molten bauxite ( $\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$ ) using electrolysis. Cryolite ( $\text{AlF}_3$ ) added to the ore in order to lower the melting point required and thus the energy required by the process.

- i. Write an half equation to show how aluminium metal is produced from the ore. [01]
- iv. What mass of aluminium metal would be produced if a current of 30,000A is applied to a cell for 1 hour. [04]
- v. In the molten mixture there is a mixture of anions which mostly consists of  $\text{O}_2^-$  and  $\text{F}^-$ . Write an equation to show which of these anions will be oxidised in the cell? [01]

**Question two (20 Marks)**

- a) State Kohlrausch's law. [01]
- b) The table below shows limiting molar conductivities of common anions and cations at 298K;

Cation	Limiting molar conductivity ( $\Lambda^\circ_m$ ) $\text{Scm}^2\text{mol}^{-1}$	Anion	Limiting molar conductivity ( $\Lambda^\circ_m$ ) $\text{Scm}^2\text{mol}^{-1}$
$\text{H}^+$	349.6	$\text{OH}^-$	199.1
$\text{Na}^+$	50.1	$\text{Cl}^-$	76.3
$\text{K}^+$	73.5	$\text{Br}^-$	78.1
$\text{Ca}^{2+}$	119.0	$\text{Ac}^-$	40.0
$\text{Mg}^{2+}$	106.0	$\text{SO}_4^{2-}$	160.0

Use it to answer the questions (a) and (b) that follow.

Explain the difference in conductivity between;

- i.  $\text{H}^+$  ion and  $\text{Na}^+$  ion
  - ii.  $\text{Na}^+$  and  $\text{Ca}^{2+}$
  - iii.  $\text{SO}_4^{2-}$  and  $\text{OH}^-$  [03]
  - iv. Calculate the value of ( $\Lambda^\circ_m$ ) for  $\text{CaCl}_2$  and  $\text{MgSO}_4$  solutions. [02]
- c) The limiting molar conductivity ( $\Lambda^\circ_m$ ) values for  $\text{NaCl}$ ,  $\text{HCl}$  and  $\text{NaAc}$  are  $126.4 \text{ Scm}^2\text{mol}^{-1}$ ,  $425.9 \text{ Scm}^2\text{mol}^{-1}$ , and  $91.05 \text{ Scm}^2\text{mol}^{-1}$ , respectively. Use these values to calculate the value of ( $\Lambda^\circ_m$ ) for  $\text{HAc}$ . [03]
  - d) The conductivity of  $0.001028 \text{ mol.L}^{-1}$  acetic acid is  $4.95 \times 10^{-5} \text{ scm}^{-1}$ . Calculate its dissociation constant if ( $\Lambda^\circ_m$ )<sub>acetic acid</sub> is  $390.05 \text{ Scm}^2\text{mol}^{-1}$  [03]


- e) The conductivity of sodium chloride at 298K was determined at different concentrations and the results tabulated as below.

Concentration (mol.m <sup>-3</sup> )	Conductivity k x 10 <sup>2</sup> (Sm <sup>2</sup> )	Molar conductivity ( $\Lambda_m$ )Scm <sup>2</sup> mol <sup>-1</sup>	C <sup>1/2</sup>
0.001	1.237		
0.010	11.85		
0.020	23.15		
0.050	55.58		
0.100	106.7		

- Calculate ( $\Lambda_m$ ) for all values of concentration given, and draw a graph between ( $\Lambda_m$ ) and C<sup>1/2</sup>. Use the graph to find the value of  $\Lambda_m^\circ$ . [05]
- f) Explain how using Kohlraush law one can determine  $\Lambda_m^\circ$  for distilled water. [03]

### Question three (20 Marks)

- b) The Nickel Cadmium cell, also called an alkaline cell has the overall spontaneous reaction
- $$\text{Cd}_{(s)} + \text{NiO}(\text{OH})_{(s)} \rightarrow \text{Cd}(\text{OH})_{2(s)} + \text{Ni}(\text{OH})_{2(s)}$$
- iv. Write the equation of the reaction at the **anode**. [02]
- v. Write the equation of the reaction at the **Cathode**. [02]
- vi. Explain how the alkaline cell recharges itself. [01]
- vii. State **three** disadvantages and **two** disadvantages of the alkaline accumulator over the Lead-acid cell. [05]
- c) The diagram below shows a galvanic cell operating under standard conditions. The cell reaction taking place when the cell is functioning is:
- $$6\text{Cl}^- (\text{aq}) + 2\text{Au}^{3+} (\text{aq}) \rightarrow 3\text{Cl}_2(\text{g}) + 2\text{Au}(\text{s})$$
- With switch is open, the initial reading on the voltmeter is 0.14 V.



Write down the:

- i. NAME or FORMULA of the oxidising agent [01]
- ii. Half-reaction which takes place at the anode [02]
- iii. Cell notation for this cell [02]
- iv. Calculate the standard reduction potential of Au. [03]
- v. Switch S is now closed and the bulb lights up. How will the reading on the voltmeter now compare to the INITIAL reading of 0,14 V?. Give a reason for the answer. [02]

**Question four (20 Marks)**

- a) Calculate the Gibb's free energy in Joules for a Zn/Cu cell whose standard cell potential is +1.10V. Explain this cell is spontaneous? [02]
- b) Use the standard reduction potentials below to calculate the equilibrium constant for the following reaction at 25°C. [03]
 
$$3\text{I}_2(\text{s}) + 2\text{Al}(\text{s}) \rightarrow 6\text{I}^-(\text{aq}) + 2\text{Al}^{3+}(\text{aq})$$

$$\text{I}_2(\text{s}) + 2\text{e}^- \rightarrow 2\text{I}^-(\text{aq}) \quad E_0 = +0.54\text{V} \dots\dots\dots(\text{i})$$

$$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s}) \quad E_0 = -1.66\text{V} \dots\dots\dots(\text{ii})$$
- c) A concentration cell is made using two Zn half cells, one with  $[\text{Zn}^{2+}(\text{aq})] = 0.1\text{M}$  and the other  $[\text{Zn}^{2+}(\text{aq})] = 1.0\text{M}$ . if  $\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s}) \quad E_0 = -0.76\text{V}$ , Calculate the potential developed across the terminals of this cell. Comment on the cell potential when the two half cells have the same concentration. [04]
- d)
  - i. Differentiate the following terms; Resistivity and Conductivity [02]
  - ii. Calculate the resistance of a 12m copper wire whose diameter is 0.01m ( $\rho_{\text{Cu}} = 1.68 \times 10^{-8}\text{m}$ ) [02]
- e) Differentiate between metallic conduction and electrolytic conduction. [03]

- f) The resistance of a conductivity cell filled with  $0.1 \text{ mol.L}^{-1}$  KCl is  $100 \Omega$ . If the resistance of the same cell when filled with  $0.02 \text{ mol.L}^{-1}$  solution is  $520 \Omega$ . Calculate conductivity and molar conductivity of the  $0.02 \text{ mol.L}^{-1}$  KCl solution. The resistivity ( $\rho$ ) of  $0.1 \text{ mol.L}^{-1}$  KCl solution is  $1.29 \text{ sm}^{-1}$ . [04]

**Question five (20 Marks)**

- a) What is Potentiometric Titration? [02]  
b) Explain the principle in Potentiometric titration. [02]  
c) Describe the method of Potentiometric titration. [04]  
d) Name FOUR types of Potentiometric titrations. Give a brief description of each of these types of titration. [04]  
e) What is the main advantage of potentiometric titration? [02]  
f) How do you determine the endpoint of this titration? [02]  
g) Mention one use of quinhydrone? [02]  
h) Which electrode is used as a reference electrode? [02]