

22



KIBABII UNIVERSITY

UNIVERSITY EXAMINATIONS
(MAIN EXAMINATIONS)
2020/2021 ACADEMIC YEAR

FIRST YEAR SECOND SEMESTER EXAMINATIONS
FOR THE
DEGREE OF BACHELOR OF SCIENCE COMPUTER SCIENCE

COURSE CODE: MAT 212

COURSE TITLE: LINEAR ALGEBRA

DATE: 22/7/2021

TIME: 2 PM – 4 PM

INSTRUCTIONS TO CANDIDATE

- Answer question ONE and any other TWO questions

TIME: 2 HOURS

This Paper Consists of 4 Printed Pages. Please Turn Over

QUESTION ONE [30 marks]

- a) Let V be a vector space over a field F . given that $U = \{u_1, u_2, \dots, u_n\}$ is a subset of V , explain the meaning of the following; U is
- a subspace of V (2 mks)
 - linearly independent (2 mks)
 - a basis for V (2 mks)

b) consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 2 & 1 & 0 \end{pmatrix}$$

Find the reduced row echelon equivalence of A . (3 mks)

c) solve the system

$$x_2 + x_3 - 2x_4 = -3$$

$$x_1 + 2x_2 - x_3 = 2$$

$$2x_1 + 4x_2 + x_3 - 3x_4 = -2$$

$$x_1 - 4x_2 - 7x_3 - x_4 = -19 \quad (10 \text{ mks})$$

d) find the kernel of the linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ represented by

$$T(x_1, x_2) = (x_1 - 2x_2, 0, -x_1) \quad (5 \text{ mks})$$

e) Let $T: \mathbb{R}^5 \rightarrow \mathbb{R}^7$ be a linear transformation

- Find the dimension of the kernel of T if the dimension of the range is 2 (2 mks)
- Find the rank of T if the nullity of t is 4 (2 mks)
- Find the rank of T if $\ker(T) = \{0\}$ (2 mks).

QUESTION TWO (20 marks)

- (a) If W is a nonempty subset of a vector space V , then W is a subspace of V iff the following closure conditions hold.

If u and v are in W , then $u+v$ is in W .

If u is in W and c is a scalar, then cu is in W . Prove (6 mks)

- (b) Let W be the set of all 2×2 symmetric matrices. Show that W is a subspace of the vector space $M_{2,2}$, with the standard operations of matrix addition and scalar multiplication. (4 mks)
- (c) If V and W are both subspaces of a vector space U , then the intersection of V and W is also a subspace of U . (6 mks)
- (d) Show that the subset of \mathbb{R}^2 consisting of all points on the unit circle $x^2+y^2 = 1$ is not a subspace. (4 mks)

QUESTION THREE (20 marks)

- a. Consider the set of vectors $V = \{[x, y, z]: ax + by + cz = 0\}$ where a, b, c are scalars. Show that V is a vector space. (10 mks)
- b. Let V be a vector space, then
- c. $\alpha \cdot 0 = 0$ for every scalar α
- d. $0 \cdot x = 0$ for every x in V
- e. If $\alpha \cdot x = 0$ then $\alpha = 0$ or $x = 0$ (10 mks)

QUESTION FOUR (20 marks)

- a. Show that the set of vectors

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \text{ spans } \mathbb{R}^3$$

(5 mks)

- b. If A and B are invertible matrices of size n , then AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$ (6 mks)
- c. Determine whether the set of vectors in P_2 is linearly independent or linearly dependent $S = \{1+x-2x^2, 2+5x-x^2, x+x^2\}$ (9 mks)

QUESTION FIVE (20 marks)

a. When is an $n \times n$ matrix A invertible? (2 mks)

b. If A is an invertible matrix, then its inverse is unique. Prove (7 mks)

c. Show that B is the inverse of A where,

$$A = \begin{pmatrix} -1 & 2 \\ -1 & 1 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 1 & -2 \\ 1 & -1 \end{pmatrix} \quad (5 \text{ mks})$$

d. Compute A^{-1} in two ways and show that the results are equal given that

$$A = \begin{pmatrix} 1 & 1 \\ 2 & 4 \end{pmatrix} \quad (6 \text{ mks})$$