



(Knowledge for Development)

KIBABII UNIVERSITY

UNIVERSITY EXAMINATIONS
2021/2022 ACADEMIC YEAR
SECOND YEAR SUPPLEMENTARY
EXAMINATION

FOR THE DEGREE OF BACHELOR OF SCIENCE

COURSE CODE: STA 212

COURSE TITLE: SAMPLE SURVEYS II

DATE: FRI 20/07/2022

TIME: 8:00 AM - 10:00 AM

INSTRUCTIONS TO CANDIDATES

Answer Question One and Any other TWO Questions

TIME: 2 Hours

This Paper Consists of 3 Printed Pages. Please Turn Over.

QUESTION 1: (30 marks)

(a) By use of suitable examples, describe the following sampling techniques:

1) Genealogy sampling		(2 marks)
ii)	Snowball sampling	(2 marks
" D	11	(0 1

(b) Let the sample arithmetic mean $\overline{y} = \frac{1}{n} \sum_{i=1}^{n} y_i$ be an estimator of the population

mean $\overline{Y} = \frac{1}{N} \sum_{i=1}^{N} Y_i$. Verify that \overline{y} is an unbiased estimator of \overline{Y} under:

(c) Describe stratified sampling (6 marks)

(d) Given the following data

Stratum, h	N_h	Sh
1	45	10
2	20	19
3	65	5

For a fixed sample size, n= 60, obtain n_h under the,

(i)	Optimum allocation scheme	(4 marks)
(ii)	Proportional allocation scheme	 (4 marks)

QUESTION 2: (20 marks)

(a) The sample size is to be determined such that the variance of \bar{y} should not exceed a given value, say V. In this case we find n such that $Var(\bar{y}) \leq V$. Proceed to justify that for large N, the smallest value of the sample size, $n_{smallest}$ is

$$n_{\text{smallest}} = n_{\text{e}} \text{ and } n \ge n_{\text{e}}, \text{ where } n_{\text{e}} = \frac{S^2}{v}$$
 (10 marks)

(b) Suppose we want to determine sample size, n such that the following requirement is satisfied,

$$P[I\overline{y} - \overline{Y}I \le e] = 1 - \alpha$$

Where e is the absolute estimation error and α the level of significance.

Assume
$$\overline{y}$$
 follows $N\left(\overline{Y}, \frac{N-n}{Nn}S^2\right)$. Show that $n = \left(\frac{Z_{\frac{\alpha}{2}}}{e}S\right)$ for large N. (10 marks)

QUESTION 3: (20 marks)

(i) What do you understand by the term, Power of a statistical test?

(ii) An investigator is planning a clinical trial to evaluate the efficacy of a new drug designed to reduce systolic blood pressure. The plan is to enrol participants and to them to receive either the new drug or a placebo. Systolic blood pressures will be a similar trials, the investigator expects that 10% of all participants will be a similar trials.

will drop out of the study. If the new drug shows a 5 unit reduction in mean systolic blood pressure, this would represent a clinically meaningful reduction. At 5% level of significance, how many patients should be enrolled in the trial to ensure that the power of the test is 80% to detect this difference?

(16 marks)

QUESTION 4: (20 marks)

- (a) Consider the estimation of \bar{y} under SRSWOR and SRSWR. Which of these two sampling schemes is more efficient in carrying out the estimation? (4 marks)
- (b) Suppose it is desired that the coefficient of variation, CV of \bar{y} should not exceed a given or pre-specified value of coefficient of variation, say C_0 , then the required sample size n is to be determined such that,

$$CV(\overline{y}) \le C_0 \text{ or } \frac{\sqrt{\text{var}(\overline{y})}}{\overline{Y}} \le C_0$$

Under these conditions, show that the smallest possible sample size $n_{smallest}$ is given by

$$n_{smallest} = \frac{C^2}{C_0^2}$$
, where C is the population coefficient of variation (10 marks)

(c) Describe Cluster sampling. How does it differ from Stratified sampling? (6 marks)

QUESTION 5: (20 marks)

- (a) State the advantages of systematic over simple random sampling scheme. (4 marks)
- (b) Distinguish linear systematic sampling (LSS) from Circular systematic sampling (CSS) (6 marks)
- (c) Prove that in LSS the population variance is the sum of the variations within the samples and between the samples (10 marks)