

On Generation of Measurable Covers for Measurable Sets Using Multiple Integral of Functions

Otanga Levi Olwamba¹, Aywa Shem Omukunda¹, Owino Maurice Oduor²

1. Department of Mathematics, Masinde Muliro University of Science and Technology P.O. Box 190-50100, Kakamega, Kenya.

2. Department of Mathematics and Computer Science, University of Kabianga P.O Box 2030-20200, Kericho, Kenya.

Abstract

In this article, we formulate an n -dimensional structure of measurable covers for measurable sets. Properties such as monotonicity, countable additivity and σ -finiteness of the projective tensor product of vector measure duality are largely applied.

Keywords: Measurable cover, Multiple integral, Vector measure duality.

Introduction

Over the years, vector measure integration has been used as a tool for analysis of structural properties of Banach space functions. Analysts have applied measurability concepts of the Lebesgue measure to construct measurable covers for certain subsets of the real line. In this paper, we apply multiple integral of vector valued functions with respect to projective tensor product of vector measure duality to generate measurable covers for measurable sets in the space \mathfrak{R}^n .

Basic Concepts

1. Projective Tensor Product Vector Measure Duality

Let X_1, \dots, X_n and Z be real Banach spaces with $\Phi : \Pi_{i=1}^n X_i \rightarrow Z$ being a continuous linear function. If $\mu_1 : \mathcal{R}_1 \rightarrow X_1, \dots, \mu_n : \mathcal{R}_n \rightarrow X_n$ are countably additive vector measures, then the product

Corresponding author: Owino Maurice Oduor, Department of Mathematics and Computer Science, University of Kabianga P.O Box 2030-20200, Kericho, Kenya. E-mail: morricearaka@yahoo.com.

$\Pi_{i=1}^n \mu_i$ is a countably additive vector measure defined on the ring $\Pi_{i=1}^n R_i$ generated by sets of the form $E_1 \times \dots \times E_n$. Let $G(R_1), \dots, G(R_n)$ be a set of σ -rings generated by rings R_1, \dots, R_n respectively.

If the extension of the vector measure $\Pi_{i=1}^n \mu_i : \Pi_{i=1}^n R_i \rightarrow \Pi_{i=1}^n X_i$ to a vector measure $\Pi_{i=1}^n \mu_i^* : \Pi_{i=1}^n G(R_i) \rightarrow \Pi_{i=1}^n X_i$ coincide with respect to a linear function $\Phi : \Pi_{i=1}^n X_i \rightarrow Z$, where $Z = \Phi(\mu_1^*(E_1), \dots, \mu_n^*(E_n))$ for $\mu_i^*(E_i) \in X_i$, $1 \leq i \leq n$ and $\Pi_{i=1}^n X_i$ is a Banach space, then $(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i), z' \rangle$ is called the projective tensor product of vector measure duality between Z and its dual space Z' .

2. Integrable Functions

An $\Pi_{i=1}^n G(R_i)$ -measurable function f is said to be integrable with respect to the projective tensor product of vector measure duality $(\Pi_{i=1}^n)_\Phi \langle \mu_i^*, z' \rangle$ if for every set $\Pi_{i=1}^n E_i = ((e_1, \dots, e_n) :$

$f(e_1, \dots, e_n) \neq 0) \in \Pi_{i=1}^n G(R_i)$ there exists an element $\int_n \dots \int_1 f \delta \Pi_{i=1}^n \mu_i^* \in \Pi_{i=1}^n X_i$, such that

$$\langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(E_i)}(f), Z' \rangle = \int_n \dots \int_1 f \delta (\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i), z' \rangle$$

where Z' is the dual space of $Z = \Phi(\mu_1^*(E_1), \dots, \mu_n^*(E_n))$, $\mu_i^*(E_i) \in X_i$ for $1 \leq i \leq n$ and $z' \in Z'$,

$T : L^n(\lambda) \rightarrow \Pi_{i=1}^n X_i$ is the operator given by $T(f) = \int_n \dots \int_1 f \delta_\lambda$ for $\lambda = \mu_1^* \times \dots \times \mu_n^*$ and $f \in L^n(\lambda)$.

3. Measurable Covers

Let f be an integrable function with respect to the projective tensor product of vector measure duality $(\Pi_{i=1}^n)_\Phi \langle \mu_i^*, z' \rangle$. Suppose $\Pi_{i=1}^n E_i$ and $\Pi_{i=1}^n C_i$ are measurable sets with respect to the ring $\Pi_{i=1}^n R_i$ and σ -ring $\Pi_{i=1}^n G(R_i)$ respectively. We say that $\Pi_{i=1}^n C_i = ((c_1, \dots, c_n) : f(c_1, \dots, c_n) \neq 0)$ is a measurable cover of $\Pi_{i=1}^n E_i$ in symbols $\Pi_{i=1}^n E_i \Theta \Pi_{i=1}^n C_i$, if the following conditions are satisfied; (i) $E_i \subset C_i$ for

$i = 1, \dots, n$ (ii) if $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_i), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(C_i - E_i), z' >$ then

$A_i \downarrow \emptyset$ for each $i = 1, \dots, n$. Hence, $\langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(A_i)}(f), Z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_i), z' \rangle = 0$.

Applying the property of directed projective tensor product of vector measure duality on integrable functions, (see [1,2,12]), we write $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' \rangle = LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(C_i), z' \rangle$

4. Measurable Cover Estimate Technique

Let $\Pi_{i=1}^n \mu_i^* : \Pi_{i=1}^n G(R_i) \rightarrow \Pi_{i=1}^n X_i$ be a product vector measure.

Suppose $\Pi_{i=1}^n C_i = ((c_1, \dots, c_n) : f(c_1, \dots, c_n) \neq 0)$ is a set in a σ -ring $\Pi_{i=1}^n G(R_i)$ generated by a ring $\Pi_{i=1}^n R_i$. If $\Pi_{i=1}^n E_i$ is a set in $\Pi_{i=1}^n R_i$ such that $E_i \subset C_i$ for $1 \leq i \leq n$, the set $\Pi_{i=1}^n C_i$ is said to be the best approximation of a measurable cover for a measurable set $\Pi_{i=1}^n E_i$, if given a real number $\varepsilon > 0$, there exists a measurable set $\Pi_{i=1}^n A_i$, such that

$$\int_n \dots \int_1 f \delta \Pi_{i=1}^n < \mu_i^*(A_i), z' \rangle \leq \int_n \dots \int_1 f \delta \Pi_{i=1}^n < \mu_i^*(C_i - E_i), z' \rangle \text{ and}$$

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_i), z' \rangle < \varepsilon$$

where f is an integrable function with respect to projective tensor product of vector measure duality $(\Pi_{i=1}^n)_\Phi < \mu_i^*, z' \rangle$ and z' is an element in the dual space Z' of Z (see [10]). In this case, the measure of $\Pi_{i=1}^n A_i$ gives the optimal error between the set $\Pi_{i=1}^n E_i$ and its cover $\Pi_{i=1}^n C_i$.

5. σ -finite Projective Tensor Product Vector Measure

A projective tensor product vector measure $\Pi_{i=1}^n \mu_i^*$ on a ring $\Pi_{i=1}^n R_i$ is said to be σ -finite if given any $\Pi_{i=1}^n E_i = N(f) \in \Pi_{i=1}^n R_i$, there exists a sequence $(\Pi_{i=1}^n E_{k_i})_{k=1}^\infty$ in $\Pi_{i=1}^n R_i$ such that $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' \rangle \leq \sum_{k=1}^\infty \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_{k_i}), z' \rangle$ and $\mu_i^*(E_{k_i}) < \infty$ for $1 \leq i \leq n$

Proposition 1. Let f be an integrable function with respect to $(\Pi_{i=1}^n)_\Phi < \mu_i^*, z' \rangle$ such that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{i^*}^*(A_i), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{i^*}^*(E_i), z' >$$

$$\leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{i^*}^*(C_i), z' >$$

where E_i and C_i are sets in R_i and $G(R_i)$ respectively for $1 \leq i \leq n$. If A_i is measurably covered by C_i for $1 \leq i \leq n$, then $\Pi_{i=1}^n A_i \Theta \Pi_{i=1}^n E_i$,

Proof. Let $\Pi_{i=1}^n C_i = ((c_1, \dots, c_n) : f(c_1, \dots, c_n) \neq 0) \in \Pi_{i=1}^n G(R_i)$. From the hypothesis, we have

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i - A_i), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(C_i - A_i), z' >$$

Since A_i is measurably covered by C_i for $1 \leq i \leq n$, it follows that $< T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(E_i - A_i)}(f), Z' > = 0$.

By integrability of f (see [4,5]), we have $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i - A_i), z' > = 0$. Suppose

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_{i_k}), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i - A_i), z' >, \text{ where } A_{i_k} \neq A_i \text{ for}$$

$i_k \neq i < T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(A_{i_k})}(f), Z' > = 0$. As is the case with the integral operator T_f with respect to projective

tensor product of vector measure duality (see[9, p. 485]). So $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_{i_k}), z' > = 0$

$$\Rightarrow \Pi_{i=1}^n A_i \Theta \Pi_{i=1}^n E_i.$$

Proposition 2. Let $\Pi_{i=1}^n C_i = ((c_1, \dots, c_n) : f(c_1, \dots, c_n) \neq 0)$ If a set $(\Pi_{i=1}^n E_i)$ is measurably covered by

$$(\Pi_{i=1}^n C_i) \text{ such that } \Pi_{i=1}^n E_{i_k} \uparrow \Pi_{i=1}^n E_i \text{ and } \Pi_{i=1}^n C_{i_k} \uparrow \Pi_{i=1}^n C_i, \text{ then } \cup_{k=1}^\infty \Pi_{i=1}^n E_{i_k} \Theta \cup_{k=1}^\infty \Pi_{i=1}^n C_{i_k}$$

Proof. Since $\Pi_{i=1}^n E_{i_k} \uparrow \Pi_{i=1}^n E_i$ and $\Pi_{i=1}^n C_{i_k} \uparrow \Pi_{i=1}^n C_i$, it shown in [8], that

$$\cup_{k=1}^\infty \Pi_{i=1}^n E_{i_k} = \Pi_{i=1}^n E_i \text{ and } \cup_{k=1}^\infty \Pi_{i=1}^n C_{i_k} = \Pi_{i=1}^n C_i. \text{ Let } \cup_{k=1}^\infty \Pi_{i=1}^n A_{i_k} \text{ be a measurable set with respect to}$$

$\Pi_{i=1}^n G(R_i)$ such that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(\cup_{k=1}^\infty A_{i_k}), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(\cup_{k=1}^\infty (C_{i_k} - E_{i_k})), z' >.$$

It is shown in [12] that each integrable function f with respect to vector measure duality can be identified with

the operator T . Therefore, we need to show that $\Sigma_{k=1}^{\infty} \langle T_{(\Pi_i^n)_\Phi \mu_i^*(A_{i_k})}(f), Z' \rangle = 0$.

$$\begin{aligned} & \Sigma_{k=1}^{\infty} \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(A_{i_k}), z' \rangle \geq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_{i_k} - \cup_{k=1}^{\infty} E_{i_k}), z' \rangle \\ & \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(\cup_{k=1}^{\infty} (C_{i_k} - E_{i_k})), z' \rangle. \end{aligned}$$

Under countable additivity, the result in [13] establishes that

$$\begin{aligned} & \Sigma_{k=1}^{\infty} \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(A_{i_k}), z' \rangle \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(\cup_{k=1}^{\infty} (C_{i_k} - E_{i_k})), z' \rangle \\ & \leq \Sigma_{k=1}^{\infty} \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*((C_{i_k} - E_{i_k})), z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_i - E_i), z' \rangle \end{aligned}$$

By hypothesis $\Pi_{i=1}^n E_i \ominus \Pi_{i=1}^n C_i$. So,

$$\begin{aligned} & \Sigma_{k=1}^{\infty} \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(A_{i_k}), z' \rangle = 0 < T_{(\Pi_i^n)_\Phi \mu_i^*(\cup_{k=1}^{\infty} A_{i_k})}(f), Z' \rangle = 0 \\ & \Rightarrow \Sigma_{k=1}^{\infty} \langle T_{(\Pi_i^n)_\Phi \mu_i^*(A_{i_k})}(f), Z' \rangle = 0 \end{aligned}$$

Proposition 3. If $\Pi_{i=1}^n A_i \ominus \Pi_{i=1}^n E_i$ and $\Pi_{i=1}^n A_i \ominus \Pi_{i=1}^n C_i = N(f)$, such that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \Delta C_i), z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*((E_i - C_i) \cup (C_i - E_i)), z' \rangle, \text{ then}$$

$$\langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(E_i \Delta C_i)}(f), Z' \rangle = 0 \text{ and therefore,}$$

$$\begin{aligned} & \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i), z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_i), z' \rangle \\ & = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cup C_i), z' \rangle \\ & = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cap C_i), z' \rangle. \end{aligned}$$

Proof. Since $\Pi_{i=1}^n A_i \ominus \Pi_{i=1}^n E_i$ and $\Pi_{i=1}^n A_i \ominus \Pi_{i=1}^n C_i$, by theorem on monotone classes (see [7] p.13), it follows

that

$$\begin{aligned} & \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(A_i), z' \rangle \geq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cap C_i), z' \rangle \\ & \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i), z' \rangle. \end{aligned}$$

Therefore,

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i - (E_i \cap C_i)), z' \rangle \geq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i - A_i), z' \rangle.$$

Since $\Pi_{i=1}^n A_i \ominus \Pi_{i=1}^n E_i$, then by hypothesis,

$$\begin{aligned} & \langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(E_i)}(f), Z' \rangle - \langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(E_i \cap C_i)}(f), Z' \rangle \\ & = \langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(E_i - E_i \cap C_i)}(f), Z' \rangle = 0 \end{aligned} \quad (\text{see [10], Th.14, p.485})$$

It follows that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i - E_i \cap C_i), z' \rangle = 0$$

so that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i - C_i), z' \rangle = 0.$$

Similarly, since

$$\begin{aligned} & \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(A_i), z' \rangle \geq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cap C_i), z' \rangle \\ & \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^* C_i, z' \rangle. \end{aligned}$$

$$\text{So } \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_i - (E_i \cap C_i)), z' \rangle \geq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_i - A_i), z' \rangle.$$

$$\text{By hypothesis, } \Pi_{i=1}^n A_i \ominus \Pi_{i=1}^n C_i \Rightarrow \langle T_{(\Pi_{i=1}^n)_\Phi \mu_i^*(C_i - E_i \cap C_i)}(f), Z' \rangle = 0.$$

As a consequence of the integral operator T acting on the integrable function f , (see [11], p. 34), we obtain

$$\begin{aligned} & \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_i - E_i \cap C_i), z' \rangle = 0 \Rightarrow \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(C_i - E_i), z' \rangle \\ & = 0 \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \Delta C_i), z' \rangle = 0. \end{aligned}$$

Now, consider the relation

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cup C_i), z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*((E_i \Delta C_i) \cup (E_i \cap C_i)), z' \rangle.$$

Applying results obtained in [6, 11] on vector measure additivity and the symmetric difference of measurable sets (see [3], p.3), we obtain

$$\begin{aligned} & \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cup C_i), z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \Delta C_i), z' \rangle \\ & + \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cap C_i), z' \rangle = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cup C_i), z' \rangle \\ & = \int_n \dots \int_1 \delta(\Pi_{i=1}^n)_\Phi \langle \mu_i^*(E_i \cap C_i), z' \rangle. \end{aligned}$$

Since $\Pi_{i=1}^n (E_i \cap C_i) \subset \Pi_{i=1}^n E_i \subset \Pi_{i=1}^n (E_i \cup C_i)$ and

$$\begin{aligned}
& \Pi_{i=1}^n (E_i \cap C_i) \subset \Pi_{i=1}^n C_i \subset \Pi_{i=1}^n (E_i \cup C_i) \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i \cap C_i), z' > \\
& = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > \\
& = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(C_i), z' > \\
& = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i \cup C_i), z' >
\end{aligned}$$

Proposition 4. Let $\Pi_{i=1}^n E_i = ((e_1, \dots, e_n) : f((e_1, \dots, e_n)) \neq 0)$ be a measurable set with respect to $\Pi_{i=1}^n R_i$. If

$$\begin{aligned}
& \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > < \infty, \text{ there exists a set } \Pi_{i=1}^n C_i \in \Pi_{i=1}^n G(R_i) \text{ such that} \\
& \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > = LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(C_i), z' >
\end{aligned}$$

Proof. Choose a set $(A_i, i = 1, 2, \dots, n)$ in $\Pi_{i=1}^n G(R_i)$ such that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_i), z' > N(f) = \Pi_{i=1}^n E_i.$$

by hypothesis, $\chi_{\Pi_{i=1}^n E_i} f \leq f$ for $f > 0 \Rightarrow \chi_{\Pi_{i=1}^n E_i} f \leq \chi_{\Pi_{i=1}^n A_i} f$. On applying the directed projective tensor product of vector measure duality, we obtain

$$LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(A_i), z' > \downarrow \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > \text{ for each } k \text{ (see [7], p.20), and}$$

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(A_i), z' > < \infty. \text{ Let } \Pi_{i=1}^n C_i = \bigcap_{k=1}^\infty \Pi_{i=1}^n A_{k_i} \text{ such that } \bigcap_{k=1}^\infty A_{k_i} = A_i \Rightarrow$$

$$\Pi_{i=1}^n A_i \downarrow \Pi_{i=1}^n C_i. \text{ ([7], p.20) and ([3], p. 92)}$$

$$LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(A_i), z' > \downarrow LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(C_i), z' > \text{ for each } k,$$

$$LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(C_i), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' >$$

Proposition 5. Every measurable set $\Pi_{i=1}^n E_i \in \Pi_{i=1}^n R_i$ with σ -finite projective tensor product vector

measure has a measurable cover i.e. there exists a set $N(f) = \Pi_{i=1}^n C_i$ in $\Pi_{i=1}^n G(R_i)$ such that $\int_n \dots \int_1 f$

$$\delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > = LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_{k_i}^*(C_i), z' >$$

Proof. By σ -finiteness of the projective tensor product vector measure, there exists a sequence $(A_{k_i})_{k=1}^{\infty}$ in

$\Pi_{i=1}^n R_i$ such that $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} A_{k_i}), z' > < \infty$ for all k and

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(E_i), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} A_{k_i}), z' > .$$

Suppose

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} A_{k_i}), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(E_i \cap (\cup_{k=1}^{\infty} A_{k_i})), z' >$$

$$\Rightarrow \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(E_i), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} A_{k_i}), z' > .$$

From Proposition 4, there exists a set $\Pi_{i=1}^n H_{k_i}$ in $\Pi_{i=1}^n G(R_i)$ such that $\Pi_{i=1}^n A_{k_i} \Theta \Pi_{i=1}^n H_{k_i}$. Define

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(C_i), z' > = \sum_{k=1}^{\infty} \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(H_{k_i}), z' > .$$

From the results above, we obtain $\Pi_{i=1}^n E_i \Theta \Pi_{i=1}^n C_i$. Since $E_i \subset C_i$ for $1 \leq i \leq n$ hold (see [2]). Now,

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(E_i), z' > = LUB_k \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(C_i), z' >$$

Theorem 1. If $\Pi_{i=1}^n E_i = N(f)$ is a $\Pi_{i=1}^n R_i$ - measurable set of σ -finite projective tensor product vector

measure, there exists a set $\Pi_{i=1}^n C_i$ in $\Pi_{i=1}^n G(R_i)$ such that $\Pi_{i=1}^n E_i \Theta \Pi_{i=1}^n C_i$ and

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(C_i - E_i), z' > = 0$$

Proof. By σ -finiteness of the projective tensor product of vector measure duality, there exists a sequence

$(A_{k_i})_{k=1}^{\infty}$ in $\Pi_{i=1}^n R_i$ such that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(E_i), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} A_{k_i}), z' >$$

and $< T_{(\Pi_{i=1}^n)_{\Phi} \mu_i^*(A_{k_i})}(f), Z' > < \infty$ for all k . For each k , choose a set $\Pi_{i=1}^n H_i$ in $\Pi_{i=1}^n G(R_i)$ such that

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(A_{k_i}), z' > \leq \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(H_i), z' > .$$

Since $N(f) = \Pi_{i=1}^n E_i$, it follows that $\chi_{\Pi_{i=1}^n H_i} f \leq f$ for $f > 0$. Define

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(A_{k_i}), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_{\Phi} < \mu_i^*(E_i \cap H_i), z' > .$$

Then,

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(E_i), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(\cup_{k=1}^\infty A_{k_i}), z' > \\ = \sum_{k=1}^\infty \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_{k_i}), z' > .$$

Since $\Pi_{i=1}^n E_i$ is $\Pi_{i=1}^n R_i$ -measurable, the sequence of sets $(A_{k_i})_{k=1}^\infty$ are $\Pi_{i=1}^n R_i$ -measurable. Therefore,

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(A_{k_i}), z' > < \infty .$$

By proposition 4, there exists a set $\Pi_{i=1}^n H_{k_i}$ in $\Pi_{i=1}^n G(R_i)$ such that $\Pi_{i=1}^n A_{k_i} \ominus \Pi_{i=1}^n H_{k_i}$. Define

$$\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(C_i), z' > = \sum_{k=1}^\infty \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(H_{k_i}), z' > .$$

From the results above, we have $\Pi_{i=1}^n E_i \ominus \Pi_{i=1}^n C_i$.

So, $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(C_i - E_i), z' > = \int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(\cup_{k=1}^\infty (H_{k_i} - A_{k_i})), z' >$. Since,

$\Pi_{i=1}^n A_{k_i} \ominus \Pi_{i=1}^n H_{k_i}$, it follows that $\int_n \dots \int_1 f \delta(\Pi_{i=1}^n)_\Phi < \mu_i^*(C_i - E_i), z' > = 0$.

References

- [1]. Berberian Khazag Sterling (1965), Measure and Integration, Chelsea Publishing Company Bronx, Newyork, 1-312.
- [2]. Bruckner B. and Thomson S.(1997), Real Analysis (1st ed.), Prentice Hall,1-713.
- [3]. Charalambos D. Aliprantis and Owen Burkinshaw (1989), Principles of Real Analysis (2nd ed.), Department of Mathematical sciences , Indiana University at Indianapolis, Indiana, 1-285.
- [4]. Campo Del R., Fernandez A., Ferrrando I., Mayoral F. and Naranjo F. (2010), Compactness of multiplication operators on spaces of integrable functions with respect to a vector measure, in vector measure, integration and related topics, Birkhauser Verlag, Basel Vol.201, pp.109-113.
- [5]. Curbera G.P. (1995), Banach space properties of L' of vector measure, Proc.Amer. Math. Soc. 123, 3797-3806.
- [6]. Dorlas T.C. (2010), Reminder notes for the course on measures on topological spaces, Dublin Institute for Advanced Studies, Dublin, Ireland,10,1-22.
- [7]. Elias M. and Rami Shakarchi (2005), Real Analysis (3rd.Edition), Princeton University Press, 1-48.
- [8]. Garret Birkoff (1995), Lattice theory, Colloquium Publications 25 (3rd ed),American Mathematical Society, p. 95.
- [9]. Gomez Palacio P., Lopez Molina J.A. and Rivera M.J.(2001), Some applications of Lattice finite representability in spaces of measurable functions,Turk J. Math. 25, 475-490.
- [10]. Jimenez Fernandez E. and Sanchez Perez E.A.(2012), Lattice copies of in of a Vector Measure and Strongly

Orthogonal Sequences, Journal of function spaces and applications, Hindawi Publishing Corporation, 1-15.

- [11]. Pratulananda Das and Md Mamun Ar Rashid (2002), A new sparse set topology, Vietnam Journal of Mathematics, 30, 113-120.
- [12]. Sanchez Perez E.A. (2001), Compactness arguments for spaces of p-integrable functions with respect to a vector measure and factorization of operators through Lebesgue-Bochner spaces, Illinois J. Math. 45, 907-923.
- [13]. Yaogan Mensah (2013), Facts about the Fourier-Stieltjes transform of vector measures on Compact Groups, International Journal of Analysis and Applications, 19-25.