



KIBABII UNIVERSITY

**UNIVERSITY EXAMINATIONS
2021/2022 ACADEMIC YEAR**

**FOURTH YEAR FIRST SEMESTER
SPECIAL/SUPPLEMENTARY EXAMINATIONS**

FOR THE DEGREE OF BSC (PHYSICS)

COURSE CODE: SPC412

COURSE TITLE: SOLID STATE PHYSICS II

DATE: 17/11/2022

TIME: 11:00AM-1:00PM

INSTRUCTIONS TO CANDIDATES


TIME: 2 Hours

Answer question ONE and any TWO of the remaining

KIBU observes ZERO tolerance to examination cheating

Question One

- Other than its high resistance, explain how germanium can be identified as semiconductors from silver metal. (2 mks)
- State one disadvantage of intrinsic semiconductors. (1 mk)
- Differentiate between P-type and N-type superconductors. (2 marks)
- Describe the effect of temperature on the conductivity of a semiconductor (3 marks)
- Find the conductivity and resistivity of a pure silicon crystal at temperature 300°K. The density of electron hole pair per cc at 300°K for a pure silicon crystal is 1.072×10^{10} and the mobility of electron $\mu_n = 1350 \text{ cm}^2/\text{volt-sec}$ and hole mobility $\mu_h = 480 \text{ cm}^2/\text{volt-sec}$. (3mks)
- How much donor impurity should be added to pure germanium so that its resistivity drops to 10% of its original value? Determine n and p in a p-type germanium sample whose resistivity is 0.01 ohm-cm, and also find n and p in a N-type silicon sample whose resistivity is 10 Ωcm . Resistivity of pure Ge is 44.6 Ωcm . (4 marks)
- Figure below shows the plot of log of resistivity versus reciprocal of temperature for two different semiconductors A and B. Assume that mobility is proportional to $T^{-3/2}$,



- Which material has wider band gap and why? (4mks)
- Define the Hall effect. (2 marks)
- State any four uses of Hall effect. (2 marks).
- The Hall coefficient of a specimen of doped silicon is found to be $3.66 \times 10^{-4} \text{ m}^3/\text{°C}$. The resistivity of the specimen is $8.93 \times 10^{-3} \Omega\text{m}$, assuming single carrier concentration find the mobility and density of charge carrier. (3 marks)
- Define the following terms as used in superconductivity
 - Critical temperature (1 mark)
 - Energy gap (1 marks)
- Differentiate between Type I and type II superconductors. (2 marks)

Question Two

- a) If N_c and N_v denote the density of states in the conduction band and density of states in the valence band, respectively while E_{fi} , E_{fn} and E_{fp} denote energies associated with the Fermi levels in intrinsic, N-type and P-type semiconductors, respectively, show that

$$E_{fn} - E_{fi} = kT \ln \frac{n}{n_i}$$

Show a diagrammatic representation of this equation. (10 marks)

- b) The conductivity of an intrinsic semiconductor $\sigma_i = n_i e (\mu_n + \mu_p)$ is given by

$$\sigma_i = n_i e (\mu_n + \mu_p)$$

Proceeding from this equation, show that conductivity of intrinsic semiconductor varies exponentially with increase in temperature. (10 marks)

Question Three

- a) Starting with the current density $J_{xe} = en\mu_e E_x$ along the x-axis of a rectangular slab, with e as the electron charge, n as the electron number density, μ_e as the electron mobility and E_x as the electric field along the x-axis, show that the Hall coefficient $R_H = \frac{e(n\mu_e^2 - p\mu_h^2)}{\sigma^2}$ where μ_h is the hole mobility and σ is the electric conductivity of the semiconductor. (10 Marks)
- b) Using a well labeled diagram, show how the Hall coefficient can be determined from the measurable quantities in a laboratory. Define each quantity clearly and how it can be measured. (10 marks)

Question Four

- a) Discuss the characteristic properties of superconductors under the following headlines (10 marks)
- Resistivity
 - Meissner Effect
 - Energy gap
 - Coherence length
- b) Derive:
- The London's first equation (3 marks)
 - The London's second equation (4 marks)
 - The London's penetration depth (5 marks)

Question Five

- a) Show that the dielectric constant for a system of two parallel plates of charge density $+\sigma$ and $-\sigma$ separated a distance d by a vacuum of permittivity ϵ_0 , is

$$\epsilon_r = 1 + \chi_E$$

Where χ_E is the electric susceptibility. (10mks)

- b) Discuss point defects in crystalline solids. (10mks)