



Analysis of Fecal Coliform Levels at Watering Points along the Upper Reaches of River Isiukhu in Kakamega County, Kenya

Sarah Kindiki^{1*}, Mario Kollenberg¹, Donald Siamba², Anthony Sifuna¹ and Clabe Wekesa¹

¹*Department of Biological Sciences, Masinde Muliro University of Science and Technology, P.O.Box 190-50100, Kakamega, Kenya.*

²*Department of Agriculture and Veterinary Sciences, Kibabii University, P.O.Box 1699-50200, Bungoma, Kenya.*

Authors' contributions

This work was carried out in collaboration between all authors. Author SK designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors MK and DS managed the analyses of the study. Author CW managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMB/2018/39522

Editor(s):

- (1) Foluso O. Osunsanmi, Department of Biochemistry and Microbiology, University of Zululand, South Africa.
(2) Pongsak Rattanachaikunsopon, Professor, Department of Biological Science, Faculty of Science, Ubon Ratchathani University, Thailand.

Reviewers:

- (1) O. I. Afolami, Federal University of Technology, Nigeria.
(2) Azizur Rahman, University of Toronto, Canada.
(3) Fábio Henrique Portella Corrêa de Oliveira, Universidade Federal Rural de Pernambuco, Brazil.
(4) Takashi Furukawa, School of Allied Health Sciences, Kitasato University, Japan.
(5) Ahmed Elsadek Fakh, Zagazig University, Egypt.

Complete Peer review History: <http://www.sciencedomain.org/review-history/24309>

Original Research Article

Received 4th January 2018
Accepted 13th March 2018
Published 24th April 2018

ABSTRACT

Diarrheal diseases often attributable to poor sanitary conditions and fecal contamination of drinking water remain a leading cause of mortality for children younger than five years. Water contaminated with human faeces, for example from municipal sewage, septic tanks and latrines, is of particular concern. Animal faeces also contain microorganisms that can cause diarrhea. Kakamega County in Kenya has a diarrhea prevalence rate of 20.2%, which is the highest in the country; a good proportion of these cases are believed to be water borne. This study was designed to determine

*Corresponding author: E-mail: skindiki@gmail.com;

fecal coliform levels in water samples collected from watering points along the upper reaches of River Isiukhu in Kakamega County, Kenya. Fifty-four samples were collected between August and October 2015 from nine sampling points, comprising springs and watering points along the river. Water samples were filtered on nitrocellulose filters by vacuum filtration; fecal coliform counts were conducted using membrane filters cultured on mFC agar to establish contamination levels. The results indicated that counts ranged from 200 cfu/100 ml – 1450 cfu/100 ml in river sampling points and ranged from 0 cfu/100 ml – 44 cfu/100 ml in springs sampling points. The fecal coliform counts for River Isiukhu and most springs surrounding it exceeded the WHO recommended drinking water coliform (or *E. coli*) count value of 0 cfu/100 ml indicating that water from the upper reaches of River Isiukhu and springs is not fit for drinking before treatment, especially during the wet seasons, based on WHO drinking water standards.

Keywords: Fecal-coliform; contamination; river-water; spring-water; season.

1. INTRODUCTION

Water has been classified as a natural resource and is important in sustaining life [1] reported that the accessibility and availability of clean drinking water not only plays a vital role in economic development and social welfare, but is also an important component in health, food production and poverty alleviation. Despite its significance, [2] revealed that safe potable water is not accessible by about 1.1 billion people in the world, and the hourly toll from biological contamination of drinking water is 400 deaths of children below the age of five. In most developing countries, Kenya being one of them, the demand for clean drinking water supply is growing rapidly [3]. In addition, a small percentage of people in these countries access piped water. Therefore, those who do not have access to safe drinking water, as well as those who have access but cannot afford it; rely on other sources of water of questionable quality [4, 5]. It is often assumed that spring water emerges from the ground clean and free of contaminants, especially in rural areas where industrial contamination is not present [6]. Many rural Kenyans know that drinking untreated water from surface streams and rivers is not safe but, they often assume that water emerging from the ground at a spring is clean and safe to drink.

Although not absolute, several pathogenic microorganisms have been recommended as indices of fecal pollution and act as indicators of microbiological quality of domestic water [7, 8,9].

Besides anthropogenic activities, natural phenomena are also believed to be contributing to the reduction of water quality worldwide. For example, increasingly unpredictable seasonal discharge of storm-waters into lotic ecosystems due to climate variability synergistically

compounds river discharges and the water quality [10]. Exhaustion of the natural soil nutrients has forced farmers to use organic and inorganic fertilizers in agro-ecosystems leading to high influx of run-off waters, rich in nutrients in rivers and streams. This has, in many occasions resulted into eutrophication which has both health and ecological repercussions in aquatic ecosystems.

According to [2], drinking water should contain zero fecal coliform and coliform organisms per 100 ml. However, human activities, particularly urbanization, and waste disposal and agricultural practices have greatly increased inputs of microbial and other pollutants to terrestrial and aquatic habitats [11]. Therefore, the present study sought to establish the level of fecal coliform in sampling points in the upper reaches of River Isiukhu.

The study was carried out in Kakamega County (Latitude: 0° 16' 60.00" N Longitude: 34° 45' 0.00" E). The average annual rainfall of Kakamega County is 1800mm per annum and is bimodally distributed with peaks in April-May and August-September. The driest months are from December-February. Temperatures range from a minimum of 10.3°C to a maximum of 30.8°C with an average of 20.5°C. The area is covered by Kakamega phonolites and also tertiary volcanic rocks-olivine basalts and nepheline. Kakamega County has a population of 1.66 million people and an area of 3,033.8km². The Isiukhu River originates from Nandi hills and flows through Kakamega forest, a tropical rain forest. River Isiukhu drains land use, land cover of forest, sugarcane plantation, mixed agriculture and periurban from upstream to downstream before joining River Nzoia which empties into Lake Victoria. The river flows in an east-west direction passing about 2 km from Kakamega town in the

south side. Anthropogenic activities are done along the river such as agriculture, livestock watering, laundry and bathing and sewage disposal.

2. MATERIALS AND METHODS

2.1 Sample Collection

The sampling sites were River Isiukhu and springs along it. The sites were randomly chosen and they represented the upper reaches of the

river. In the study, samples were collected in three successive months (August, September and October). Approximately 250ml of water samples were aseptically collected in duplicates from watering points using sterile containers and immediately stored in an ice box before being transported to the laboratory for analysis. Water samples were collected from a depth of about 10-20 cm below the water surface at each site to avoid potential contamination from surface water.




Fig. 1. Sampling sites (S1-S5) and (SP1-SP4) in the upper reaches of River Isiukhu Kakamega County, Kenya. Map drawn to scale using Arc GIS software (units in UTM)

2.2 Research Design

The present study adopted randomized sampling design, where Fifty four duplicate samples were collected between the month of August and October 2015 from nine sampling points along the river and springs that represented the upper reaches of River Isiukhu. The forest area (S1) provides reference data, land cover of forest, human activities like deforestation. Confluence of Lianila stream and River Isiukhu (S2) assess bacterial load of Lianila stream that drains the treated sewage effluent and farm lands. Activities such as fishing, bathing, livestock watering, and laundry are also done. Savona resort area (S3) quantifies the bacterial contribution from the recreation area.

Amalemba area (S4) is highly populated; determines the bacterial contribution of the small scale mixed agriculture and periurban area activities eg waste disposal of the households, municipal wastes, hotel and dispensaries wastes etc bathing, livestock watering, laundry are done. The drainage of this region is skewed. Waste water entry point (S5) establishes the bacterial load from treated sewer waters from municipal sewage treatment plant in the area and the slaughter house drains. All springs (SP1- SP4) are located along River Isiukhu and the study is to establish their bacterial contamination and any variation in season's i.e during dry and wet season. Contamination due to pollution likely from sewage as well as solid waste discharge during run- offs.

2.3 Determination of Fecal Coliform Levels at Watering Points and Springs along the Upper reaches of River Isiukhu

Fecal coliform levels in water samples were determined by the membrane filter procedure using mFC agar [12]. The samples collected from the river were diluted to 1:100. The biomass from both spring and the diluted river-water samples

(100 ml) was concentrated via subsequent filtration on nitrocellulose filters, 47 mm diameter with pore size of 0.45 µm by vacuum filtration. The vacuum pressure for filtration was between 50 mm to 70 mm Hg in order to avoid rupture of bacterial cells that has been observed at pressures above 80 mm Hg [13]. Following the method recommended by the American Public Health Association in 2006 [14,15] filters were then aseptically placed with grid side up onto the surface of the plates of mFC. All plates were incubated inverted in watertight plastic bags submerged in a 44.5°C water bath for 22-24 hours. Fecal coliform colonies that were observed in any shade of blue color were counted using a Quebec colony counter and recorded as colony-forming units per 100ml. The formula used was:

$$CFU/100mL = \frac{\text{Colonies counted}}{\text{mL filtered}} \times 100$$

2.4 Data Analysis

Data obtained on coliform counts was subjected to one-way ANOVA followed by Turkey's post hoc test at 95% confidence level using Winks software version 7.

3. RESULTS

Table 1 shows the mean counts of fecal coliforms from five different sampling points in River Isiukhu for three successive months. There was a significantly high ($p < 0.05$) fecal coliform concentration between sampling point, S5 and all other sampling points in the river in the month of August. Sampling point, S2 also recorded significantly higher ($p < 0.05$) coliform counts when compared to S1 in the month of October. In the month of September, S2 and S5 had visibly higher coliform counts per 100ml, though they were not significantly different from the rest of the samples. Generally, fecal coliform counts were lowest at S1 and highest at S5 in all the three month of the study.

Table 1. Means of fecal coliform count given in colony forming units /100mls for five sampling points from River Isiukhu for three consecutive months

Sample code	Environment	Aug.	Sept.	Oct.
S1	Forest (site 1)	200±0.00 ^a	600±141.42 ^a	600±424.26 ^a
S2	Confluence of Lianila stream and River Isiukhu	300±141.42 ^{ab}	1050±1343.50 ^a	1700±424.26 ^b
S3	Savona resort area	550±494.97 ^{ab}	750±636.40 ^a	950±353.55 ^{ab}
S4	Amalemba area	450±353.55 ^{ab}	900±848.53 ^a	1050±636.40 ^{ab}
S5	Waste water entry point	1450±212.13 ^c	1050±353.55 ^a	950±212.13 ^{ab}

*Means followed by the same letter within the same column are not significantly different at $P < 0.05$

Table 2. Means of Fecal coliform count given in colony forming units /100ml for four sampling points from spring for six sessions

Sample code	Environment	Aug.	Sept.	Oct.
SP1	Confluence of Lianila stream and River Isiukhu	0±0.00 ^a	5±7.07 ^a	6.5±4.95 ^a
SP2	Savona resort area	3±1.41 ^a	3±2.83 ^a	44±46.67 ^a
SP3	Amalemba area	2.5±3.54 ^a	6.5±4.95 ^a	6.5±4.95 ^a
SP4	Waste water entry point	5.5±4.95 ^a	6.5±2.12 ^a	36.5±43.13 ^a

**Means followed by the same letter within the same column are not significantly different at P <0.05*

3.1 Fecal Coliform Count for Four Sampling Points from Springs along River Isiukhu

Table 2 shows the mean counts for fecal coliform from four different springs for three consecutive months. Generally, the month of October recorded higher coliform count compared to September and August. The month of August recorded few counts across all the springs, in the three months studied. However, it is important to note that there was no significant difference in the counts for all the months studied. In the month of August, spring SP1 recorded no coliform counts, however, increasing counts were recorded in the month of September and October.

4. DISCUSSION

Generally, coliforms are the most common group of indicator organisms used in water quality monitoring [16]. Furthermore, Alotaibi [17] elaborated that the presence of fecal coliform indicates the presence of potential fecal contamination and the presence of possible pathogenic microorganisms and to determine the health risk to the consumers. In the present study, significant variation of fecal coliform counts in River Isiukhu was reported during the three months of the study. This difference is attributed to rainfall variation within the three months. Lower rainfalls are normally experienced in the month of August than in the month of October. Similar observations have been made elsewhere, for instance, Wolf [18] reported that in the dry season, there were fewer incidences of fecal pollution in the water supply system but high fecal contamination of drinking water increases during wet season.

Results in the present study further indicated significant differences in means for fecal coliforms in samples taken from the different sampling points in River Isiukhu. This finding

agrees with a previous study carried out on River Danube which showed lower bacterial pollution on upstream of the River and higher levels of fecal pollution in the middle part and downstream of the River [19]. However, the finding in this study disagrees with that of Uwimpuhwe [20] where total and fecal coliform from different sampling points in River Nyabarongo showed insignificant differences. This difference could be attributed to the fact that the current study of River Isiukhu included survey of sampling points with different anthropogenic activities taking place at every point.

According to the WHO guideline values for bacteriological parameters, the total and fecal coliform bacteria should be 0 cfu/100ml in water intended for drinking. However, in this study the fecal coliform counts for River Isiukhu, exceeded the WHO recommended drinking water guideline value. These results are supported by previous studies conducted in rural areas [9,21]. The studies found that microbiological parameters counts for river water in rural areas were above the permissible limits and were a potential hazard to public health [21]. Even though, WHO recognizes that these targets would be difficult to achieve in some cases, especially in rural communities with untreated water provisions, and recommends that in these settings, the guidelines values would be seen as goals for the future, but not an immediate prerequisite [22].

Although findings from the study reported insignificant difference in fecal coliform counts from different springs during the three months studied, the water was unsafe to drink based on the World Health Organization drinking water standards [23]. Both protected and unprotected springs had bacterial counts in excess of the WHO standard, suggesting that water treatment from all sources is necessary to ensure their cleanliness and safety. The spring water contamination could be due to what Narain Rai and Sharma [24] termed as lack of sanitation or improper waste disposal. The researchers further

explained that 40% or more of the disease outbreaks were attributed to consumption of polluted ground water. Furthermore, presence of coliforms in drinking water sources indicates need for treatment and proper sanitation which is necessary for drinking [25]. Further the destruction of microbial ecosystems through deforestation, high spring water temperatures, averaging 24.4°C, may be contributing to the observed bacterial abundance.

Findings from the study indicate that springs have more fecal coliform contaminants during wet season. This finding is not different from that of Ofoma [26] where spring water was highly contaminated with fecal coliform during wet seasons compared to dry seasons. They suggested that the contamination was due to pollution; most likely from sewage as well as solid waste discharge during run-offs. This is non-point sources typically wet-weather where they diffuse in nature, in that they do not enter water bodies from any single point (e.g. urban litter, contaminated refuse, domestic pet/wildlife excrement and failing sewer lines). This suggests that emphasis on points of use (POU) treatment methods, decontamination of protected and unprotected springs, and behavioral interventions to improve sanitation practices are needed.

5. CONCLUSION

Results obtained from the study indicated significant contamination of River and spring water at different sites. The levels of fecal coliforms were higher than the accepted levels. In addition, higher fecal coliform levels were recorded during wet than dry seasons, concluding that water from the upper reaches of River Isiukhu and spring is not fit for drinking before treatment especially during wet seasons based on the WHO drinking water standards and the water quality. It is therefore recommended that water from both the stream and springs in the upper reaches of River Isiukhu should be treated before use.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ashbolt NJ, Grabow WO, Snozzi M. Indicators of microbial water quality. IWA Publishing. 2001;289-316.

2. WHO. Microbial aspects. In Guidelines for drinking water quality. World Health Organisation 3rd ed. Geneva. 2006;1.
3. Gelover S, Gomez LA, Reyes K, Leal MT. A practical demonstration of water disinfection using TiO₂ films and sunlight. Water Res. 2006;40(17):3274-3280.
4. Gadgil A. Drinking water in developing countries. Annual Review of Energy and the Environment Resources. 1998;23(1):253-286.
5. Odonkor ST, Ampofo JK. Escherichia coli as an indicator of bacteriological quality of water: An overview. Microbiology Research. 2013;4(1):2.
6. Wampler PJ, Sisson AJ. Spring flow, bacterial contamination, and water resources in rural Haiti. Environmental Earth Sciences. 2011;62(8):1619-1628. Available:<http://doi.org/10.1007/s12665-010-0645-9>
7. Kabler PW, Clark HF. Coliform group and fecal coliform organisms as indicators of pollution in drinking water. J Am Water Works Assoc. 1960;52(12):1577-1579.
8. World Health Organization. Water, sanitation and health. Geneva, World Health Organization. (2006). Guidelines for drinking-water quality: First addendum to volume 1, Recommendations. World Health Organization. 2003;1.
9. Abera S, Zeyinudin A, Kebede B, Deribew A, Ali S, Zemene E. Bacteriological analysis of drinking water sources. Afr. J. Microbiol. Res. 2011;5(18):2638-2641.
10. Onyando ZA, Shivoga WA, Lung'ayia H, Ochieno DW, Agevi H, Kigen C. The influence of land use on nutrient regime in a tropical stream. Elixir Pollut. 2013; 2013(64):19290-19294.
11. Smol JP. Pollution of lakes and rivers: A paleoenvironmental perspective. John Wiley & Sons; 2009.
12. Dufour AP, Strickland ER, Cabelli VJ. Membrane filter method for enumerating *Escherichia coli*. Applied and Environmental Microbiology. 1981;41(5):1152-1158.
13. Kepner RL, Pratt JR. Use of fluorochromes for direct enumeration of total bacteria in environmental samples. past and present. Microbiol Rev. 1994;58(4):603-615.
14. Wehr HM, Frank JF, American Public Health Association (Eds.). Standard methods for the examination of dairy

- products. Washington, DC: American Public Health Association. 2004;327-404.
15. Britton LJ, Greeson PE. Methods for collection and analysis of aquatic biological and microbiological samples. U.S. Geological Survey Techniques of Water Resources Investigations. 1987;5(A4):37-40.
 16. Sibanda T, Chigor VN, Okoh AI. Environ Monit Assess. 2013;185:6579. Available:<https://doi.org/10.1007/s10661-012-3048-4>
 17. Alotaibi EL. Bacteriological assessment of urban water sources in Khamis Mushait Governorate, Southwestern Saudi Arabia. Int J Health Geogr. 2009;8(1): 1.
 18. Wolf AT. The transboundary freshwater dispute database project. WATER INT. 1999;24(2):160-163.
 19. Kavka GG, Kasimir GD, Farnleitner AH. Microbiological water quality of the River Danube (km 2581-km 15). Longitudinal variation of pollution as determined by standard parameters. Eigenverlag ICPDR. 2006;415–421.
 20. Uwimpuhwe M. Microbiological assessment of water quality and prevalence of waterborne diseases in rural areas of Masaka, Rwanda (Doctoral Dissertation); 2014.
 21. Chigor VN, Umoh VJ, Okuofu CA, Ameh JB, Igbinosa EO, Okoh AI. Water quality assessment. Surface water sources used for drinking and irrigation in Zaria, Nigeria are a public health hazard. Environ. Monit. Assess. 2012;184(5):3389-3400.
 22. WHO/UNICEF Joint Water Supply, Sanitation Monitoring Programme, World Health Organization, UNICEF. Progress on drinking water and sanitation: Special focus on sanitation. World Health Organization; 2008.
 23. World Health Organization. Progress on sanitation and drinking water: 2015 update and MDG assessment. World Health Organization; 2015.
 24. Narain Rai JP, Sharma HC. Bacterial Contamination of Ground Water in Rural Areas of North Uttarpradesh. Indian Journal Environmental Health. 1995; 37(1):37-41.
 25. Christine L, Moe, Richard D, Rheingans. Global challenges in water, sanitation and health. Journal of Water and Health. 04.IWA Publishing; 2006.
 26. Ofoma AE, Omologbe DA, Aigberua P. Physico-chemical quality of groundwater in parts of Port Harcourt City, Eastern Niger Delta, Nigeria. Water Resour. Res. 2005; 16:18-24.

© 2018 Kindiki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/24309>